

Mean Sum Square Prime Labeling of Some Snake Graphs

Sunoj B S*, Mathew Varkey T K

Department of Mathematics, Government Polytechnic College, Attingal, Kerala, India

ABSTRACT

Mean sum square prime labeling of a graph is the labeling of the vertices with $\{0,1,2,\dots,p-1\}$ and the edges with mean of the square of the sum of the labels of the incident vertices or mean of the square of the sum of the labels of the incident vertices and one, depending on the sum is even or odd. The greatest common incidence number of a vertex (**gcin**) of degree greater than one is defined as the greatest common divisor of the labels of the incident edges. If the **gcin** of each vertex of degree greater than one is one, then the graph admits mean sum square prime labeling. Here we identify some snake graphs for mean sum square prime labeling.

Keywords : Graph Labeling, Sum Square, Greatest Common Incidence Number, Prime Labeling, Snake Graphs.

I. INTRODUCTION

All graphs in this paper are simple, finite and undirected. The symbol $V(G)$ and $E(G)$ denotes the vertex set and edge set of a graph G . The graph whose cardinality of the vertex set is called the order of G , denoted by p and the cardinality of the edge set is called the size of the graph G , denoted by q . A graph with p vertices and q edges is called a (p,q) -graph.

A graph labeling is an assignment of integers to the vertices or edges. Some basic notations and definitions are taken from [2],[3] and [4]. Some basic concepts are taken from [1] and [2]. In [5], we introduced the concept of sum square prime labeling and proved the result for some cycle related graphs. In [6], [7], [8], [9], we proved the result for some path related graphs, some snake related graphs, some tree graphs, triangular belt, jelly fish graph, some star related graphs. In this paper we introduced mean sum square prime labeling using the concept greatest common incidence number of a vertex. We proved that some snake graphs admit mean sum square prime labeling.

Definition: 1.1 Let G be a graph with p vertices and q edges. The greatest common incidence number (**gcin**)

of a vertex of degree greater than or equal to 2, is the greatest common divisor (gcd) of the labels of the incident edges.

II. MAIN RESULTS

Definition 2.1 Let $G = (V, E)$ be a graph with p vertices and q edges. Define a bijection

$f : V(G) \rightarrow \{0,1,2,3,\dots,p-1\}$ by $f(v_i) = i-1, 1 \leq i \leq p$.

Define a 1-1 mapping

$f_{mssp}^* : E(G) \rightarrow$ set of natural numbers N by

$$f_{mssp}^*(uv) = \frac{\{f(u)+f(v)\}^2}{2}, \text{ when } f(u)+f(v) \text{ is even.}$$

$$f_{mssp}^*(uv) = \frac{\{f(u)+f(v)\}^2+1}{2}, \text{ when } f(u)+f(v) \text{ is odd.}$$

The induced function f_{mssp}^* is said to be a mean sum square prime labeling, if the **gcin** of each vertex of degree at least 2, is 1.

Definition 2.2 A graph which admits mean sum square prime labeling is called a mean sum square prime graph.

Theorem 2.1 Triangular snake T_n admits mean sum square prime labeling.

Proof: Let $G = T_n$ and let $v_1, v_2, \dots, v_{2n-1}$ are the vertices of G

Here $|V(G)| = 2n-1$ and $|E(G)| = 3n-3$

Define a function $f : V \rightarrow \{0,1,2,3,\dots,2n-2\}$ by

$$f(v_i) = i-1, i = 1, 2, \dots, 2n-1.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{mssp}^* is defined as follows

$$f_{mssp}^*(v_i v_{i+1}) = 2i^2 - 2i + 1, \quad i = 1, 2, \dots, 2n-2$$

$$f_{mssp}^*(v_{2i-1} v_{2i+1}) = 8i^2 - 8i + 2, \quad i = 1, 2, \dots, n-1$$

Clearly f_{mssp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_1) &= \text{gcd of } \{f_{mssp}^*(v_1 v_2), f_{mssp}^*(v_1 v_3)\} \\ &= \text{gcd of } \{1, 2\} = 1. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{2n-1}) &= \text{gcd of } \{f_{mssp}^*(v_{2n-3} v_{2n-1}), f_{mssp}^*(v_{2n-1} v_{2n-2})\} \\ &= \text{gcd of } \{8n^2 - 20n + 13, 8n^2 - 24n + 18\} \\ &= \text{gcd of } \{4n-5, 8n^2 - 24n + 18\} \\ &= \text{gcd of } \{4n^2 - 12n + 9, 4n-5\} \\ &= \text{gcd of } \{n-1, 4n-5\} \\ &= \text{gcd of } \{n-1, n-2\} \\ &= 1 \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{i+1}) &= \text{gcd of } \{f_{mssp}^*(v_i v_{i+1}), f_{mssp}^*(v_{i+1} v_{i+2})\} \\ &= \text{gcd of } \{2i^2 - 2i + 1, 2i^2 + 2i + 1\} \\ &= \text{gcd of } \{4i, 2i^2 - 2i + 1\} \\ &= \text{gcd of } \{i, 2i^2 - 2i + 1\} \\ &= 1, \quad i = 1, 2, \dots, 2n-3. \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1. Hence T_n , admits mean sum square prime labeling.

Theorem 2.2 Pentagonal snake P_n admits mean sum square prime labeling, if $(n+1) \not\equiv 0 \pmod{5}$.

Proof: Let $G = P_n$ and let $v_1, v_2, \dots, v_{4n-3}$ are the vertices of G

$$\text{Here } |V(G)| = 4n-3 \text{ and } |E(G)| = 5n-5$$

Define a function $f: V \rightarrow \{0, 1, 2, 3, \dots, 4n-4\}$ by

$$f(v_i) = i-1, i = 1, 2, \dots, 4n-3.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{mssp}^* is defined as follows

$$f_{mssp}^*(v_i v_{i+1}) = 2i^2 - 2i + 1, \quad i = 1, 2, \dots, 4n-4$$

$$f_{mssp}^*(v_{4i-3} v_{4i+1}) = 32i^2 - 32i + 8, \quad i = 1, 2, \dots, n-1$$

Clearly f_{mssp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_1) &= \text{gcd of } \{f_{mssp}^*(v_1 v_2), f_{mssp}^*(v_1 v_4)\} \\ &= \text{gcd of } \{1, 8\} \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{4n-3}) &= \text{gcd of } \{f_{mssp}^*(v_{4n-3} v_{4n-4}), f_{mssp}^*(v_{4n-3} v_{4n-7})\} \\ &= \text{gcd of } \{32n^2 - 72n + 41, 32n^2 - 96n + 72\}, \end{aligned}$$

$$\begin{aligned} &= \text{gcd of } \{8(2n-3)^2, 32n^2 - 72n + 41\}, \\ &= \text{gcd of } \{2n-3, (2n-3)(16n-12)+5\} \\ &= \text{gcd of } \{5, 2n-3\} \\ &= 1 \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{i+1}) &= \text{gcd of } \{f_{mssp}^*(v_i v_{i+1}), f_{mssp}^*(v_{i+1} v_{i+2})\} \\ &= 1, \quad i = 1, 2, \dots, 4n-5 \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1. Hence P_n , admits mean sum square prime labeling.

Theorem 2.3 Alternate triangular snake $A(T_n)$ admits mean sum square prime labeling,

if n is odd and triangle starts from the first vertex.

Proof: Let $G = A(T_n)$ and let $v_1, v_2, \dots, v_{\frac{3n-1}{2}}$ are the vertices of G

$$\text{Here } |V(G)| = \frac{3n-1}{2} \text{ and } |E(G)| = 2n-2$$

Define a function $f: V \rightarrow \{0, 1, 2, 3, \dots, \frac{3n-3}{2}\}$ by

$$f(v_i) = i-1, i = 1, 2, \dots, \frac{3n-1}{2}.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{mssp}^* is defined as follows

$$f_{mssp}^*(v_i v_{i+1}) = 2i^2 - 2i + 1, i = 1, 2, \dots, \frac{3n-3}{2}$$

$$f_{mssp}^*(v_{3i-2} v_{3i}) = 18i^2 - 24i + 8, i = 1, 2, \dots, \frac{n-1}{2}$$

Clearly f_{mssp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_1) &= \text{gcd of } \{f_{mssp}^*(v_1 v_2), f_{mssp}^*(v_1 v_3)\} \\ &= \text{gcd of } \{1, 2\} \\ &= 1. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{i+1}) &= \text{gcd of } \{f_{mssp}^*(v_i v_{i+1}), f_{mssp}^*(v_{i+1} v_{i+2})\} \\ &= 1, \quad i = 1, 2, \dots, \frac{3n-5}{2} \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1. Hence $A(T_n)$, admits mean sum square prime labeling.

Theorem 2.4 Alternate triangular snake $A(T_n)$ admits mean sum square prime labeling,

if n is odd and triangle starts from the second vertex.

Proof: Let $G = A(T_n)$ and let $v_1, v_2, \dots, v_{\frac{3n-1}{2}}$ are the vertices of G

Here $|V(G)| = \frac{3n-1}{2}$ and $|E(G)| = 2n-2$

Define a function $f: V \rightarrow \{0, 1, 2, 3, \dots, \frac{3n-3}{2}\}$ by

$$f(v_i) = i-1, i = 1, 2, \dots, \frac{3n-1}{2}.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{mssp}^* is defined as follows

$$f_{mssp}^*(v_i v_{i+1}) = 2i^2 - 2i + 1, i = 1, 2, \dots, \frac{3n-3}{2}$$

$$f_{mssp}^*(v_{3i-1} v_{3i+1}) = 18i^2 - 12i + 2, i = 1, 2, \dots, \frac{n-1}{2}$$

Clearly f_{mssp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_{i+1}) &= \text{gcd of } \{ f_{mssp}^*(v_i v_{i+1}), \\ &\quad f_{mssp}^*(v_{i+1} v_{i+2}) \} \\ &= 1, \quad i = 1, 2, \dots, \frac{3n-5}{2} \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{\frac{3n-1}{2}}) &= \text{gcd of } \{ f_{mssp}^*(v_{\frac{3n-1}{2}} v_{\frac{3n-3}{2}}), \\ &\quad f_{mssp}^*(v_{\frac{3n-1}{2}} v_{\frac{3n-5}{2}}) \} \\ &= \text{gcd of } \left\{ \frac{9n^2 - 24n + 17}{2}, \frac{9n^2 - 30n + 25}{2} \right\}, \end{aligned}$$

put $n = 2k-1$

$$\begin{aligned} &= \text{gcd of } \{ 18k^2 - 42k + 25, 18k^2 - 48k + 32 \}, \\ &= \text{gcd of } \{ 6k - 7, 18k^2 - 48k + 32 \}, \\ &= \text{gcd of } \{ 3k - 3, 6k - 7 \}, \\ &= \text{gcd of } \{ 3k - 4, 3k - 3 \} = 1. \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1.

Hence $A(T_n)$, admits mean sum square prime labeling.

Theorem 2.5 Alternate triangular snake $A(T_n)$ admits mean sum square prime labeling,

if n is even and triangle starts from the first vertex.

Proof: Let $G = A(T_n)$ and let $v_1, v_2, \dots, v_{\frac{3n}{2}}$ are the vertices of G

Here $|V(G)| = \frac{3n}{2}$ and $|E(G)| = 2n-1$

Define a function $f: V \rightarrow \{0, 1, 2, 3, \dots, \frac{3n-2}{2}\}$ by

$$f(v_i) = i-1, i = 1, 2, \dots, \frac{3n}{2}.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{mssp}^* is defined as follows

$$f_{mssp}^*(v_i v_{i+1}) = 2i^2 - 2i + 1, i = 1, 2, \dots, \frac{3n-2}{2}$$

$$f_{mssp}^*(v_{3i-2} v_{3i}) = 18i^2 - 24i + 8, i = 1, 2, \dots, \frac{n}{2}$$

Clearly f_{mssp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_{i+1}) &= \text{gcd of } \{ f_{mssp}^*(v_i v_{i+1}), \\ &\quad f_{mssp}^*(v_{i+1} v_{i+2}) \} \\ &= 1, \quad i = 1, 2, \dots, \frac{3n-4}{2} \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{\frac{3n}{2}}) &= \text{gcd of } \{ f_{mssp}^*(v_{\frac{3n}{2}} v_{\frac{3n-2}{2}}), \\ &\quad f_{mssp}^*(v_{\frac{3n}{2}} v_{\frac{3n-4}{2}}) \} \\ &= \text{gcd of } \left\{ \frac{9n^2 - 18n + 10}{2}, \frac{9n^2 - 24n + 16}{2} \right\}, \end{aligned}$$

put $n = 2k$

$$\begin{aligned} &= \text{gcd of } \{ 18k^2 - 18k + 5, 18k^2 - 24k + 8 \}, \\ &= \text{gcd of } \{ 6k - 3, 18k^2 - 24k + 8 \}, \\ &= \text{gcd of } \{ 3k - 1, 6k - 3 \}, \\ &= \text{gcd of } \{ 3k - 2, 3k - 1 \} \\ &= 1. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_1) &= \text{gcd of } \{ f_{mssp}^*(v_1 v_2), f_{mssp}^*(v_1 v_3) \} \\ &= \text{gcd of } \{ 1, 2 \} \\ &= 1. \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1.

Hence $A(T_n)$, admits mean sum square prime labeling.

Theorem 2.6 Alternate triangular snake $A(T_n)$ admits mean sum square prime labeling,

if n is even and triangle starts from the second vertex.

Proof: Let $G = A(T_n)$ and let $v_1, v_2, \dots, v_{\frac{3n-2}{2}}$ are the vertices of G

Here $|V(G)| = \frac{3n-2}{2}$ and $|E(G)| = 2n-3$

Define a function $f: V \rightarrow \{0, 1, 2, 3, \dots, \frac{3n-4}{2}\}$ by

$$f(v_i) = i-1, i = 1, 2, \dots, \frac{3n-2}{2}.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{mssp}^* is defined as follows

$$f_{mssp}^*(v_i v_{i+1}) = 2i^2 - 2i + 1, i = 1, 2, \dots, \frac{3n-4}{2}$$

$$f_{mssp}^*(v_{3i-1} v_{3i+1}) = 18i^2 - 12i + 2, i = 1, 2, \dots, \frac{n-2}{2}$$

Clearly f_{mssp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_{i+1}) &= \text{gcd of } \{ f_{mssp}^*(v_i v_{i+1}), \\ &\quad f_{mssp}^*(v_{i+1} v_{i+2}) \} \\ &= 1, \quad i = 1, 2, \dots, \frac{3n-6}{2} \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1.

Hence $A(T_n)$, admits mean sum square prime labeling.

Theorem 2.7 Let G be the graph obtained by replacing each edge of path P_n by triangles and pentagons alternately. G admits mean sum square prime labeling, if n is odd and $(n+5) \not\equiv 0 \pmod{10}$.

Proof: Let G be the graph and let $v_1, v_2, \dots, v_{3n-2}$ are the vertices of G

Here $|V(G)| = 3n-2$ and $|E(G)| = 4n-4$

Define a function $f: V \rightarrow \{0,1,2,3, \dots, 3n-3\}$ by

$$f(v_i) = i-1, \quad i = 1, 2, \dots, 3n-2.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{mssp}^* is defined as follows

$$f_{mssp}^*(v_i v_{i+1}) = 2i^2 - 2i + 1, \quad i = 1, 2, \dots, 3n-3$$

$$f_{mssp}^*(v_{6i-5} v_{6i-3}) = 72i^2 - 120i + 50, \quad i = 1, 2, \dots, \frac{n-1}{2}$$

$$f_{mssp}^*(v_{6i-3} v_{6i+1}) = 72i^2 - 48i + 8, \quad i = 1, 2, \dots, \frac{n-1}{2}$$

Clearly f_{mssp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_1) &= \text{gcd of } \{f_{mssp}^*(v_1 v_2), f_{mssp}^*(v_1 v_3)\} \\ &= \text{gcd of } \{1, 2\} \\ &= 1. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{3n-2}) &= \text{gcd of } \{f_{mssp}^*(v_{3n-2} v_{3n-3}), \\ &\quad f_{mssp}^*(v_{3n-2} v_{3n-6})\} \\ &= \text{gcd of } \{18n^2 - 42n + 25, 18n^2 - 60n + 50\}, \\ &= \text{gcd of } \{18n^2 - 42n + 25, 2(3n-5)^2\}, \\ &= \text{gcd of } \{3n-5, (3n-5)(6n-4)+5\} \\ &= \text{gcd of } \{5, 3n-5\} = 1 \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{i+1}) &= \text{gcd of } \{f_{mssp}^*(v_i v_{i+1}), \\ &\quad f_{mssp}^*(v_{i+1} v_{i+2})\} \\ &= 1, \quad i = 1, 2, \dots, 3n-4 \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1. Hence G , admits mean sum square prime labeling.

Theorem 2.8 Let G be the graph obtained by replacing each edge of path P_n by pentagons and triangles alternately. G admits mean sum square prime labeling, if n is odd.

Proof: Let G be the graph and let $v_1, v_2, \dots, v_{3n-2}$ are the vertices of G

Here $|V(G)| = 3n-2$ and $|E(G)| = 4n-4$

Define a function $f: V \rightarrow \{0,1,2,3, \dots, 3n-3\}$ by

$$f(v_i) = i-1, \quad i = 1, 2, \dots, 3n-2.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{mssp}^* is defined as follows

$$f_{mssp}^*(v_i v_{i+1}) = 2i^2 - 2i + 1, \quad i = 1, 2, \dots, 3n-3$$

$$f_{mssp}^*(v_{6i-5} v_{6i-1}) = 72i^2 - 96i + 32, \quad i = 1, 2, \dots, \frac{n-1}{2}$$

$$f_{mssp}^*(v_{6i-1} v_{6i+1}) = 72i^2 - 24i + 2, \quad i = 1, 2, \dots, \frac{n-1}{2}$$

Clearly f_{mssp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_1) &= \text{gcd of } \{f_{mssp}^*(v_1 v_2), f_{mssp}^*(v_1 v_5)\} \\ &= \text{gcd of } \{1, 8\} \\ &= 1. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{3n-2}) &= \text{gcd of } \{f_{mssp}^*(v_{3n-2} v_{3n-3}), \\ &\quad f_{mssp}^*(v_{3n-2} v_{3n-4})\} \\ &= \text{gcd of } \{18n^2 - 42n + 25, 18n^2 - 48n + 32\}, \\ &= \text{gcd of } \{6n-7, 18n^2 - 48n + 32\}, \\ &= \text{gcd of } \{6n-7, 3n-3\}, \\ &= \text{gcd of } \{3n-4, 3n-3\} \\ &= 1 \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{i+1}) &= \text{gcd of } \{f_{mssp}^*(v_i v_{i+1}), \\ &\quad f_{mssp}^*(v_{i+1} v_{i+2})\} \\ &= 1, \quad i = 1, 2, \dots, 3n-4 \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1. Hence G admits mean sum square prime labeling.

I. REFERENCES

- [1]. Apostol. Tom M, Introduction to Analytic Number Theory, Narosa, (1998). F Harary, Graph Theory, Addison-Wesley, Reading, Mass, (1972)
- [2]. Joseph A Gallian, A Dynamic Survey of Graph Labeling, The Electronic Journal of Combinatorics(2016), #DS6, pp 1 – 408.
- [3]. T K Mathew Varkey, Some Graph Theoretic Generations Associated with Graph Labeling, PhD Thesis, University of Kerala 2000.
- [4]. Sunoj B S, Mathew Varkey T K, Sum Square Prime Labeling of Some Cycle Related Graphs, International Journal of Informative & Futuristic Research, Vol.4, Issue 10, pp 7846-7851.

- [5]. Sunoj B S, Mathew Varkey T K, Sum Square Prime Labeling of Some Path Related Graphs, International Journal of Research in Engineering and Applied Sciences, Vol.7, Issue 6, pp 58-62
- [6]. Sunoj B S, Mathew Varkey T K, Sum Square Prime Labeling of Some Tree Graphs, International Journal of Recent Scientific Research, Vol.8, Issue 6, pp 17447-17449.
- [7]. Sunoj B S, Mathew Varkey T K, Sum Square Prime Labeling of Some Snake Graphs, International Research Journal of Human Resources and Social Sciences, Vol.4, Issue 6, pp 313-319.
- [8]. Sunoj B S, Mathew Varkey T K, A Note on Sum Square Prime Labeling, Journal of Emerging Technologies and Innovative Research, Vol.4, Issue 9, pp 213-218